

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
**WO 03/013540 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/506**,  
31/519, A61P 35/02

(21) International Application Number: PCT/EP02/08941

(22) International Filing Date: 9 August 2002 (09.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/311,690 10 August 2001 (10.08.2001) US

(71) Applicant (for all designated States except AT, US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(71) Applicant (for AT only): **NOVARTIS PHARMA GMBH** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(71) Applicants (for all designated States except US): **GSF - NATIONAL RESEARCH INSTITUTE FOR ENVIRONMENT AND HEALTH** [DE/DE]; Ingolstädter Landstrasse 1, 85764 Oberschleissheim (DE). **THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 W. Seventh Street, Austin, TX 78701 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DONATO, Nicholas, J.** [US/US]; 815 Sandpiper, Sugar Land, TX 77478 (US). **FABBRO, Doriano** [CH/CH]; Im

Lee 54, CH-4144 Arlesheim (CH). **MANLEY, Paul, William** [GB/CH]; Bruggweg 12, CH-4144 Arlesheim (CH). **MESTAN, Jürgen** [DE/DE]; Zugmantelstrasse 6, 79211 Denzlingen (DE). **WARMUTH, Markus** [DE/DE]; Von-der-Tann Strasse 23, 81319 Starnberg (DE). **HALLEK, Michael** [DE/DE]; Brunnenstrasse 40, 86938 Schondorf (DE). **TALPAZ, Moshe** [US/US]; 6114 Bankside Drive, Houston, TX 77096 (US). **WU, Ji** [US/US]; 6206 Bankside Drive, Houston, TX 77096 (US).

(74) Agent: **GROS, Florent**; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NO, NZ, OM, PH, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW.

(84) Designated States (regional): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/013540 A1

(54) Title: USE OF C-SRC INHIBITORS ALONE OR IN COMBINATION WITH ST1571 FOR THE TREATMENT OF LEUKAEMIA

(57) Abstract: The invention relates to a combination which comprises (a) at least one compound decreasing the c-Src activity and (b) N-[5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl]-4-(3-pyridyl)-2-pyrimidine-amine or the monomethanesulfonate salt thereof; to pharmaceutical compositions comprising said combinations; and to a method of treating a warm-blooded animal having leukaemia, especially chronic myelogenous leukaemia, comprising administering to the animal at least one compound inhibiting the activity of a member of the Src kinase family, the Tec kinase family or a Raf kinase inhibitor, in particular inhibiting the c-Src protein tyrosine kinase activity or inhibiting simultaneously the c-Src protein tyrosine kinase activity and the Bcr-Abl tyrosine kinase activity, alone or in combination with a Bcr-Abl inhibitor, in particular N-[5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl]-4-(3-pyridyl)-2-pyrimidine-amine.

Use of c-Src inhibitors alone or in combination with STI571 for the treatment of leukaemia

The invention relates to a method of treating a warm-blooded animal, especially a human, having leukaemia comprising administering to the animal at least one compound inhibiting the c-Src protein tyrosine kinase activity, especially those compounds mentioned herein, in a quantity which is effective against leukaemia; to a method of treating a warm-blooded animal, especially a human, having leukaemia comprising administering to the animal (a) at least one compound decreasing the c-Src activity and (b) N-{5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl}-4-(3-pyridyl)-2-pyrimidine-amine (STI571) in a quantity which is jointly therapeutically effective against leukaemia; a combination which comprises (a) at least one compound decreasing the c-Src activity and (b) STI571 or the monomethanesulfonate salt thereof and optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use; a pharmaceutical composition comprising such a combination; the use of a compound inhibiting the c-Src protein tyrosine kinase activity or the use of the combination of (a) and (b) for the preparation of a medicament for the delay of progression or treatment of leukaemia; and to a commercial package or product comprising such a combination of (a) and (b) or at least one compound inhibiting the c-Src protein tyrosine kinase activity together with instructions for use thereof in the treatment of leukaemia.

Protein tyrosine kinases catalyze the phosphorylation of specific tyrosine residues. One member of this class of enzymes is the c-Src protein tyrosine kinase. Surprisingly, it has now been found that compounds inhibiting the c-Src protein tyrosine kinase activity, especially the compounds described hereinafter, are effective against leukaemia. Furthermore, it was surprisingly found that the effect in treating leukaemia of a combination which comprises (a) at least one compound decreasing the c-Src activity and (b) STI571 or the monomethanesulfonate salt thereof is greater than the effects that can be achieved with either type of combination partner alone, i.e. greater than the effects of a monotherapy using only one of the combination partners (a) and (b) as defined herein.

Hence, in a first embodiment, the present invention relates to a method of treating a warm-blooded animal having leukaemia comprising administering to the animal at least one compound inhibiting the c-Src protein tyrosine kinase activity in a quantity which is

- 2 -

therapeutically effective against leukaemia, in which method said compounds can also be present in the form of their pharmaceutically acceptable salts.

In a second embodiment, the present invention relates to a method of treating a warm-blooded animal having leukaemia comprising administering to the animal (a) at least one compound decreasing the c-Src activity and (b) STI571 in a quantity which is jointly therapeutically effective against leukaemia.



Furthermore, the present invention relates to a combination which comprises (a) at least one compound decreasing the c-Src activity and (b) STI571, wherein the active ingredients are present in each case in free form or in the form of a pharmaceutically acceptable salt, and optionally at least one pharmaceutically acceptable carrier; for simultaneous, separate or sequential use.

In a broader sense, the present invention relates to a method of treating a warm-blooded animal having leukaemia, in particular comprising administering to the animal at least one compound inhibiting the activity of a member of the Src kinase family, in particular src, yes, hck, fyn, lyn, lck, blk, fgr or Yrk, the activity of a member of the Btk or Tec kinase family or a Raf kinase inhibitor, e.g. BAY 43-9006, in a quantity which is therapeutically effective against leukaemia alone or in combination with a Bcr-Abl inhibitor, in particular STI571.


The term leukemia as used herein includes, but is not limited to, chronic myelogenous leukaemia (CML) and acute lymphocyte leukaemia (ALL), especially Philadelphia-chromosome positive acute lymphocyte leukaemia (Ph+ ALL). Preferably, the variant of leukaemia to be treated by the methods disclosed herein is CML.

The term "method of treatment" as used herein includes a treatment effecting the delay of progression of leukemia. The term "delay of progression" as used herein means in particular the administration of a medicament to patients being in a pre-stage or in an early phase of leukaemia, in which patients, for example, a pre-form or an early form of leukaemia is diagnosed or which patients are in a condition, e.g. a condition resulting from an accident, under which it is likely that a corresponding disease will develop.


The term "compounds inhibiting the c-Src protein tyrosine kinase activity" as used herein means such compounds having an  $IC_{50}$  in the range of 1 to 3000 nM, preferably in the range of 1 to 500 nM, in the proliferation test using bcr-Abl transfected 32D cells described hereinafter. The term includes, but is not limited to, compounds belonging to the structure classes of pyrrolopyrimidines, especially pyrrolo[2,3-d]pyrimidines, purines, pyrazo-pyrimidines, especially pyrazo[3,4-d]pyrimidines, pyrazopyrimidines, especially pyrazo[3,4-d]pyrimidines and pyridopyrimidines, especially pyrido[2,3-d]pyrimidines. Preferably, the term relates to those compounds disclosed in WO 96/10028, WO 97/28161, WO97/32879 and WO97/49706 and, more preferably, to the single compounds of formulae I to VIII, most preferably to the compound of formula I and V, in particular the compound of formula I.




- 4 -




(III)




(IV)




(V)



(VI)



(VII)



(VIII)

The compounds which are generically and specifically disclosed in WO 96/10028, WO 97/28161, WO97/32879 and WO97/49706, in each case in particular in the compound claims and the final products of the working examples, are hereby incorporated into the

present application by reference to these publications. The compounds can be prepared and administered as described in the cited documents, respectively. The compound of formula I can be prepared and formulated as described in WO 96/10028. The compound of formula II and its preparation is disclosed in Example 111c3 of WO 97/16452. The compound of formula IV can be prepared in analogy thereof. Both latter compounds can be formulated as described in WO 97/16452. The compound of formula III is discussed by R. Gamse et al. in J. Bone Miner. Res. 14 (Suppl. 1), 1999, S487. The compound of formula V is also known as PP1. The preparation of PP1 is described by T. Schindler, F. Sicheri et al in Molecular Cell, 1999 (3), 639, 647. The compounds of formula VI, VII and VIII are described in the following documents and the literature cited therein: J.M. Hamby et al, J. Med. Chem. 40, 1997, 2296-2303; R.L. Panek et al, J. Pharmacol. Exp. Ther. 283, 1997, 1433-1444; and S.R. Klutchko et al, J. Med. Chem. 41, 1998, 3276-3292.

STI571 can be prepared and administered as described in WO 99/03854, especially the monomesylate salt of STI571 can be formulated as described in Examples 4 and 6 of WO 99/03854. STI571 can also be administered as marketed under the trademark GLIVEC™ or GLEEVEC™.

The term "compounds decreasing the c-Src activity" as used herein includes, but is not limited to, compounds inhibiting the c-Src protein tyrosine kinase activity as defined above and to SH2 interaction inhibitors such as those disclosed in WO97/07131 and WO97/08193. Preferably, in the present invention compounds decreasing the c-Src activity are SH2 interaction inhibitors or, more preferably, compounds inhibiting the c-Src protein tyrosine kinase activity as defined above.

It will be understood that references to the pharmacologically active compounds mentioned herein are meant to also include the pharmaceutically acceptable salts. If compounds inhibiting the c-Src protein tyrosine kinase activity or a combination partner (a) or (b) have, for example, at least one basic center, they can form acid addition salts. The combination partners (a) and (b) having an acid group (for example COOH) can also form salts with bases. The pharmacologically active compounds mentioned herein may also be used in form of a hydrate or include other solvents used for crystallization. STI571, i.e. the combination partner (b), is preferably used in the present invention in the form of its monomesylate salt.

Additionally, the present invention relates to a method of treating a warm-blooded animal having leukaemia comprising administering to the animal at least one compound inhibiting the c-Src protein tyrosine kinase activity and the Bcr-Abl tyrosine kinase activity, in a quantity which is therapeutically effective against leukaemia, in which method said compounds can also be present in the form of their pharmaceutically acceptable salts. Preferably, such compound is a compound of formula V.

The utility of the compounds inhibiting the c-Src protein tyrosine kinase activity for the treatment of leukemia can be demonstrated, e.g., in the proliferation test using bcr-Abl transfected 32D cells as follows:

Bcr-Abl-transfected 32D cells (32D pGD p210 Bcr-Abl; Bazzoni, G.; et al. J. Clin. Invest. (1996), 98(2), 521-528) are cultured in RPMI 1640 (BioConcept, Allschwil, Switzerland; cat. No.: 1-41F01), 10 % fetal calf serum, 2 mM glutamine. 10000 cells in 50  $\mu$ L per well are seeded into flat bottom 96 well tissue culture plates. Complete medium alone (for controls) or serial threefold dilutions of compounds are added in triplicates to a final volume of 100  $\mu$ L and the cells are incubated at 37 °C, 5 % CO<sub>2</sub> for 65 to 72 h. The cell proliferation reagent WST-1 (Roche Diagnostics GmbH; cat.no.: 1 664 807) is added at 10  $\mu$ L per well followed by 2 h incubation at 37 °C. Colour development, depending on the amount of living cells, is measured at 440 nm. The effect for each compound is calculated as percent inhibition of the value (OD<sub>440</sub>) obtained for the control cells (100 %) and plotted against the compound concentrations. The IC<sub>50</sub>s are calculated from the dose response curves by graphic extrapolation.

Compounds inhibiting the growth of 32D-Bcr-Abl cells can be further tested on IL-3 dependent 32D wt cells to prove the specificity of the compounds for the bcr-Abl kinase and to exclude compound toxicity.

A combination which comprises (a) at least one compound decreasing the c-Src activity and (b) STI571, wherein the active ingredients are present in each case in free form or in the form of a pharmaceutically acceptable salt, and optionally at least one pharmaceutically acceptable carrier will be referred to hereinafter as a COMBINATION OF THE INVENTION.

The nature of proliferative diseases like leukemia is multifactorial. Under certain circumstances, drugs with different mechanisms of action may be combined. However, just



- 8 -

considering any combination of drugs having different mode of action does not necessarily lead to combinations with advantageous effects.

All the more surprising is the experimental finding that the administration of a COMBINATION OF THE INVENTION, results not only in a beneficial effect, especially a synergistic therapeutic effect, e.g. with regard to slowing down, arresting or reversing the progress of leukaemia or a longer duration of drug response, but also in further surprising beneficial effects, e.g. less side-effects, an improved quality of life and a decreased mortality and morbidity, compared to a monotherapy applying only one of the pharmaceutically active ingredients used in the COMBINATION OF THE INVENTION.

A further benefit is that lower doses of the active ingredients of the COMBINATION OF THE INVENTION can be used, for example, that the dosages need not only often be smaller, but can be also applied less frequently, or can be used in order to diminish the incidence of side-effects. This is in accordance with the desires and requirements of the patients to be treated.

The utility of the COMBINATION OF THE INVENTION for the treatment of leukemia can be demonstrated, e.g., in the proliferation test using bcr-Abl transfected 32D cells as follows:

The proliferation test using bcr-Abl transfected 32D cells with a COMBINATION OF THE INVENTION is carried out as described above with the following changes. Two combination partners are mixed in fixed ratios. Threefold serial dilutions of this mixture or the combination partners alone are added to the cells seeded in 96 well tissue culture plates as described above. The effects on 32D-bcr-Abl cell proliferation of a COMBINATION OF THE INVENTION is evaluated and compared with the effects of the single combination partners using CalcuSyn, a dose-effect analyzer software for single and multiple drugs (distributed by Biosoft, Cambridge).

One particular benefit of the present invention is the fact that the leukaemia that can be treated with a compound inhibiting the c-Src protein tyrosine kinase activity or with the COMBINATION OF THE INVENTION can be such leukaemia which is resistant to monotherapy employing STI571 as sole active agent, e.g. leukaemia of such patients who initially had responded to STI571 and then relapsed. Very especially, compounds inhibiting

the c-Src protein tyrosine kinase activity and COMBINATIONS OF THE INVENTION can be used for the treatment of patients in the advanced stage (blast crisis phase) of CML.

The person skilled in the pertinent art is fully enabled to select further relevant test models to prove the hereinbefore and hereinafter mentioned beneficial effects on leukaemia of a compound inhibiting the c-Src protein tyrosine kinase activity or of a COMBINATION OF THE INVENTION. The pharmacological activity of a compound inhibiting the c-Src protein tyrosine kinase activity or a COMBINATION OF THE INVENTION may, for example, be demonstrated in a suitable clinical study. Suitable clinical studies are, for example, open label non-randomized, dose escalation studies in patients with advanced leukaemia. Such studies prove in particular the synergism observed with the COMBINATIONS OF THE INVENTION. The beneficial effects on leukaemia can be determined directly through the results of these studies or by changes in the study design which are known as such to a person skilled in the art. For example, the combination partner (b) can be administered with a fixed dose and the dose of the combination partner (a) is escalated until the Maximum Tolerated Dosage is reached. Alternatively, a placebo-controlled, double blind study can be conducted in order to prove the benefits of the COMBINATION OF THE INVENTION mentioned herein.

In one embodiment of the invention, the compound inhibiting the c-Src protein tyrosine kinase activity is selected from pyrrolopyrimidines, especially pyrrolo[2,3-d]pyrimidines, purines, pyrazopyrimidines, especially pyrazo[3,4-d]pyrimidines, pyrazopyrimidines, especially pyrazo[3,4-d]pyrimidines and pyridopyrimidines, especially pyrido[2,3-d]pyrimidines. Particularly preferred are the compounds of formula I, II, III, IV, V, VI, VII and VIII, especially the compound of formula I and formula V.

Especially preferred is a combination comprising a compound of formula I and STI571 or the pharmaceutically acceptable salts thereof. Furthermore, especially preferred is a combination comprising a compound of formula V and STI571 or the pharmaceutically acceptable salts thereof.

The invention pertains also to the use of at least one compound inhibiting the c-Src protein tyrosine kinase activity or of the COMBINATION OF THE INVENTION for the treatment of leukaemia and for the preparation of a medicament for the treatment of leukaemia.

The COMBINATION OF THE INVENTION can be a combined preparation or a pharmaceutical composition.

The term "a combined preparation", as used herein defines especially a "kit of parts" in the sense that the combination partners (a) and (b) as defined above can be dosed independently or by use of different fixed combinations with distinguished amounts of the combination partners (a) and (b), i.e., simultaneously or at different time points. The parts of the kit of parts can then, e.g., be administered simultaneously or chronologically staggered, that is at different time points and with equal or different time intervals for any part of the kit of parts. Very preferably, the time intervals are chosen such that the effect on the treated disease in the combined use of the parts is larger than the effect which would be obtained by use of only any one of the combination partners (a) and (b). The ratio of the total amounts of the combination partner (a) to the combination partner (b) to be administered in the combined preparation can be varied, e.g. in order to cope with the needs of a patient sub-population to be treated or the needs of the single patient which different needs can be due to the particular disease, age, sex, body weight, etc. of the patients. Preferably, there is at least one beneficial effect, e.g., a mutual enhancing of the effect of the combination partners (a) and (b), in particular a synergism, e.g. a more than additive effect, additional advantageous effects, less side effects, a combined therapeutical effect in a non-effective dosage of one or both of the combination partners (a) and (b), and very preferably a strong synergism of the combination partners (a) and (b).

It is one objective of this invention to provide a pharmaceutical composition comprising a quantity, which is jointly therapeutically effective against leukaemia comprising the COMBINATION OF THE INVENTION. In this composition, the combination partners (a) and (b) can be administered together, one after the other or separately in one combined unit dosage form or in two separate unit dosage forms. The unit dosage form may also be a fixed combination.

The pharmaceutical compositions for separate administration of the combination partners (a) and (b) and for the administration in a fixed combination, i.e. a single galenical compositions comprising at least two combination partners (a) and (b), according to the invention can be prepared in a manner known per se and are those suitable for enteral, such as oral or rectal,

and parenteral administration to mammals (warm-blooded animals), including man, comprising a therapeutically effective amount of at least one pharmacologically active combination partner alone or in combination with one or more pharmaceutically acceptable carries, especially suitable for enteral or parenteral application.

Novel pharmaceutical composition contain, for example, from about 10 % to about 100 %, preferably from about 20 % to about 60 %, of the active ingredients. Pharmaceutical preparations for the combination therapy for enteral or parenteral administration are, for example, those in unit dosage forms, such as sugar-coated tablets, tablets, capsules or suppositories, and furthermore ampoules. If not indicated otherwise, these are prepared in a manner known per se, for example by means of conventional mixing, granulating, sugar-coating, dissolving or lyophilizing processes. It will be appreciated that the unit content of a combination partner contained in an individual dose of each dosage form need not in itself constitute an effective amount since the necessary effective amount can be reached by administration of a plurality of dosage units.

In particular, a therapeutically effective amount of each of the combination partner of the COMBINATION OF THE INVENTION may be administered simultaneously or sequentially and in any order, and the components may be administered separately or as a fixed combination. For example, the method of treatment of leukaemia according to the present invention may comprise (i) administration of the combination partner (a) in free or pharmaceutically acceptable salt form and (ii) administration of a combination partner (b) in free or pharmaceutically acceptable salt form, simultaneously or sequentially in any order, in jointly therapeutically effective amounts, preferably in synergistically effective amounts, e.g. in daily dosages corresponding to the amounts described herein. The individual combination partners of the COMBINATION OF THE INVENTION can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. Furthermore, the term administering also encompasses the use of a pro-drug of a combination partner that convert *in vivo* to the combination partner as such. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

The effective dosage of each of compounds inhibiting the c-Src protein tyrosine kinase activity and of the combination partners employed in the COMBINATION OF THE

INVENTION may vary depending on the particular compound or pharmaceutical composition employed, the mode of administration, the condition being treated, the severity of the condition being treated. Thus, the dosage regimen the COMBINATION OF THE INVENTION is selected in accordance with a variety of factors including the route of administration and the renal and hepatic function of the patient. A physician, clinician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the single active ingredients required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentration of the active ingredients within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the active ingredients' availability to target sites.


N-[5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl]-4-(3-pyridyl)-2-pyrimidine-amine monomesylate, is preferably administered to a human in a dosage in the range of about 5 to 850 mg/day, more preferably 25 to 600 mg/day and most preferably 100 to 300 mg/day. Unless stated otherwise herein, the compound is preferably administered from one to four times per day. Compounds inhibiting the c-Src protein tyrosine kinase activity, e.g. the compound of formula I, is preferably administered orally to a human in a dosage in the range of about 100 to 2000 mg/day, more preferably 500 to 1500 mg/day, e.g. 1000 mg/day.

If BAY 43-9006 is employed as a combination partner, it is preferably administered orally at doses of up to 800 mg twice daily.


Moreover, the present invention provides a commercial package comprising as active ingredients COMBINATION OF THE INVENTION, together with instructions for simultaneous, separate or sequential use thereof in the treatment of leukaemia.

What is claimed is:

1. Combination which comprises (a) at least one compound decreasing the c-Src activity protein tyrosine kinase and (b) N-{5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl}-4-(3-pyridyl)-2-pyrimidine-amine, wherein the active ingredients are present in each case in free form or in the form of a pharmaceutically acceptable salt, and optionally at least one pharmaceutically acceptable carrier; for simultaneous, separate or sequential use.
2. Combination according to claim 1 wherein compound inhibiting the c-Src protein tyrosine kinase activity is selected from a compound of formula I




and a compound of formula V




3. Combination according to claim 1 or 2 wherein compound (b) is used in the form of its monomethanesulfonate salt.
4. A combination according to any one of claims 1 to 3 for use in the therapeutic or diagnostic treatment of the animal or human body.
5. Use of a combination according to any one of claims 1 to 3 for the manufacture of a medicament for the treatment of leukaemia.
6. Method of treating a warm-blooded animal having leukaemia comprising administering to the animal at least one compound inhibiting the activity of a member of the Src kinase family, the Btk kinase family, the Tec kinase family or a Raf kinase inhibitor, in a quantity which is therapeutically effective against leukaemia, alone or in combination with a Bcr-Abl inhibitor.
7. Method of treating a warm-blooded animal having leukaemia comprising administering to the animal at least one compound inhibiting the c-Src protein tyrosine kinase activity in a quantity which is therapeutically effective against leukaemia, in which method said compounds can also be present in the form of their pharmaceutically acceptable salts.
8. Method according to claim 7, wherein the compound inhibiting the c-Src protein tyrosine kinase activity is selected from a compound of formula I

- 15 -




and a compound of formula V




9. Method of treating a warm-blooded animal having leukaemia comprising administering to the animal (a) at least one compound decreasing the c-Src activity and (b) N-{5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl}-4-(3-pyridyl)-2-pyrimidine-amine in a quantity which is jointly therapeutically effective against leukaemia.
10. Method according to claim 9, wherein the compound inhibiting the c-Src protein tyrosine kinase activity is selected from a compound of formula I



- 16 -




and a compound of formula V



11. Method of treating a warm-blooded animal having leukaemia comprising administering to the animal at least one compound inhibiting the c-Src protein tyrosine kinase activity and the Bcr-Abl tyrosine kinase activity, in a quantity which is therapeutically effective against leukaemia, in which method said compounds can also be present in the form of their pharmaceutically acceptable salts.
12. Method according to claim 11, wherein the compound inhibiting the c-Src protein tyrosine kinase activity and the Bcr-Abl tyrosine kinase activity is a compound of formula V

- 17 -



(V).

13. Method according to claim 6, 7, 9 or 11 wherein said leukaemia is resistant to monotherapy employing N-{5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl}-4-(3-pyridyl)-2-pyrimidine-amine as sole active agent.
14. Method according to claim 6, 7, 9 or 11 wherein said leukaemia is chronic myelogenous leukaemia.
15. Use of a compound inhibiting the c-Src protein tyrosine kinase activity for the manufacture of a medicament for the treatment of leukaemia.
16. Use of a compound inhibiting the c-Src protein tyrosine kinase activity and the Bcr-Abl tyrosine kinase activity for the manufacture of a medicament for the treatment of leukaemia.
17. A pharmaceutical composition comprising a quantity which is jointly therapeutically effective against leukaemia of a combination according to any one of claims 1 to 3 and at least one pharmaceutically acceptable carrier.
18. A commercial package comprising at least one c-Src protein tyrosine kinase activity inhibitor together with instructions for use thereof in the treatment of leukaemia.
19. A commercial package comprising (a) at least one compound decreasing the c-Src protein tyrosine kinase activity and (b) N-{5-[4-(4-methyl-piperazino-methyl)-benzoylamido]-2-methylphenyl}-4-(3-pyridyl)-2-pyrimidine-amine in free form or in the form of a

- 18 -

pharmaceutically acceptable salt, together with instructions for simultaneous, separate or sequential use thereof in the treatment of leukaemia.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/08941

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K31/506 A61K31/519 A61P35/02

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, EMBASE, MEDLINE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MARLEY S B ET AL: "THE TYROSINE KINASE INHIBITOR STI571, LIKE INTERFERON-ALPHA, PREFERENTIALLY REDUCES THE CAPACITY FOR AMPLIFICATION OF GRANULOCYTE-MACROPHAGE PROGENITORS FROM PATIENTS WITH CHRONIC MYELOID LEUKEMIA" EXPERIMENTAL HEMATOLOGY, NEW YORK, NY, US, vol. 28, May 2000 (2000-05), pages 551-557, XP002949104 ISSN: 0301-472X abstract  --- -/--	15, 16, 18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

12 November 2002

Date of mailing of the international search report

25/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/EP 02/08941

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DRUMMOND M W ET AL: "TYROSINE KINASE INHIBITORS IN THE TREATMENT OF CHRONIC MYELOID LEUKEMIA: SO FAR SO GOOD?" BLOOD REVIEWS, EDINBURGH, GB, vol. 15, no. 2, June 2001 (2001-06), pages 85-95, XP001036635 page 88, right-hand column, last paragraph -page 92, left-hand column</p>	15,16,18
X	<p>M. W. DRUMMOND ET AL: "TARGETING THE BCR-ABL TYROSINE KINASE IN CHRONIC MYELOID LEUKEMIA87 1" NEW ENGLAND JOURNAL OF MEDICINE, THE, MASSACHUSETTS MEDICAL SOCIETY, WALTHAM, MA, US, vol. 344, no. 14, 5 April 2001 (2001-04-05), pages 1084-1086, XP001036564 ISSN: 0028-4793 page 87, left-hand column -page 91, right-hand column</p>	15,16,18
X	<p>DRUKER B J ET AL: "LESSONS LEARNED FROM THE DEVELOPMENT OF AN ABL TYROSINE KINASE INHIBITOR FOR CHRONIC MYELOGENOUS LEUKEMIA" JOURNAL OF CLINICAL INVESTIGATION, NEW YORK, NY, US, vol. 105, no. 1, January 2000 (2000-01), pages 3-7, XP001014734 ISSN: 0021-9738 the whole document</p>	15,16,18
X	<p>AMOUI M ET AL: "SRC FAMILY-SELECTIVE TYROSINE KINASE INHIBITOR, PP1, INHIBITS BOTH FCEPSILONRI- AND THY-1-MEDIATED ACTIVATION OF RAT BASOPHILIC LEUKEMIA CELLS" EUROPEAN JOURNAL OF IMMUNOLOGY, WEINHEIM, DE, vol. 27, no. 8, August 1997 (1997-08), pages 1881-1886, XP001105330 ISSN: 0014-2980 abstract</p>	15,16,18
X	<p>DORSEY J F ET AL: "THE PYRIDO2,3-DPYRIMIDINE DERIVATIVE PD180970 INHIBITS P210BCR-ABL TYROSINE KINASE AND INDUCES APOPTOSIS OF K562 LEUKEMIC CELLS" CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, BALTIMORE, MD, US, vol. 60, 15 June 2000 (2000-06-15), pages 3127-3131, XP002949108 ISSN: 0008-5472 the whole document</p>	15,16,18

-/--

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/08941

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 10028 A (CIBA-GEIGY AG) 4 April 1996 (1996-04-04) cited in the application claims 1-14 ---	1-19
A	WO 97 16452 A (NOVARTIS AG) 9 May 1997 (1997-05-09) cited in the application claims 1-16 ---	1-19
A	HAMBY J M ET AL: "STRUCTURE-ACTIVITY RELATIONSHIPS FOR A NOVEL SERIES OF PYRIDOÄ2,3-DÜPYRIMIDINE TYROSINE KINASE INHIBITORS" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 40, no. 15, 1997, pages 2296-2303, XP002164503 ISSN: 0022-2623 cited in the application the whole document ---	1-19
A	GAMSE R ET AL: "NVP-AAK980, A NOVEL 6-PHENYLAMINO-2-ALKYLAMINO-PURINE DERIVATIVE, POTENTLY INHIBITS THE TYROSINE KINASE SRC AND BONE RESORPTION" JOURNAL OF BONE AND MINERAL RESEARCH, NEW YORK, NY, US, vol. 14, September 1999 (1999-09), page S487 XP000901380 ISSN: 0884-0431 cited in the application the whole document ---	1-19
A	KLUTCHKO S R ET AL: "2-Substituted AMINOPYRIDOÄ2,3-DÜPYRIMIDIN-7(8H)-ONES. STRUCTURE-ACTIVITY RELATIONSHIPS AGAINST SELECTED TYROSINE KINASES AND IN VITRO AND IN VIVO ANTICANCER ACTIVITY" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 41, no. 17, 1998, pages 3276-3292, XP002191992 ISSN: 0022-2623 cited in the application the whole document --- -/--	1-19

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/08941

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PANEK R L ET AL: "IN VITRO PHARMACOLOGICAL CHARACTERIZATION OF PD 166285, A NEW NANOMOLAR POTENT AND BROADLY ACTIVE PROTEIN TYROSINE KINASE INHIBITOR" JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, AMERICAN SOCIETY FOR PHARMACOLOGY AND, US, vol. 283, no. 3, 1997, pages 1433-1444, XP002191995 ISSN: 0022-3565 cited in the application the whole document</p> <p>---</p>	1-19
A	<p>HANKE J H ET AL: "DISCOVERY OF A NOVEL, POTENT, AND SRC FAMILY-SELECTIVE TYROSINE KINASE INHIBITOR" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 271, no. 2, 12 January 1996 (1996-01-12), pages 695-701, XP002051826 ISSN: 0021-9258 abstract</p> <p>---</p>	1-19
A	<p>TOPALY J ET AL: "SYNERGISTIC ACTIVITY OF THE NEW ABL-SPECIFIC TYROSINE KINASE INHIBITOR STI571 AND CHEMOTHERAPEUTIC DRUGS ON BCR-ABL-POSITIVE CHRONIC MYELOGENOUS LEUKEMIA CELLS" LEUKEMIA, MACMILLAN PRESS LTD, US, vol. 15, May 2001 (2001-05), pages 342-347, XP001039741 ISSN: 0887-6924 abstract</p> <p>---</p>	1-19
A	<p>KANO Y ET AL: "IN VITRO CYTOTOXIC EFFECTS OF A TYROSINE KINASE INHIBITOR STI571 IN COMBINATION WITH COMMONLY USED ANTILEUKEMIC AGENTS" BLOOD, W.B.SAUNDERS COMPAGNY, ORLANDO, FL, US, vol. 97, no. 7, 1 April 2001 (2001-04-01), pages 1999-2007, XP001035243 ISSN: 0006-4971 abstract</p> <p>---</p>	1-19

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/08941

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TYLER THIESING J ET AL: "EFFICACY OF STI571, AN ABL TYROSINE KINASE INHIBITOR, IN CONJUNCTION WITH OTHER ANTILEUKEMIC AGENTS AGAINST BCR-ABL-POSITIVE CELLS" BLOOD, W.B.SAUNDERS COMPAGNY, ORLANDO, FL, US, vol. 96, no. 9, 1 November 2000 (2000-11-01), pages 3195-3199, XP001035245 ISSN: 0006-4971 abstract</p> <p>-----</p>	1-19



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08941

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9610028	A	04-04-1996	AT 199553 T	15-03-2001
			AU 694801 B2	30-07-1998
			AU 3564395 A	19-04-1996
			BR 9509048 A	06-01-1998
			CA 2200210 A1	04-04-1996
			CN 1164234 A ,B	05-11-1997
			DE 69520282 D1	12-04-2001
			DE 69520282 T2	09-08-2001
			DK 783505 T3	02-07-2001
			WO 9610028 A1	04-04-1996
			EP 0783505 A1	16-07-1997
			ES 2157344 T3	16-08-2001
			FI 971225 A	14-05-1997
			GR 3035996 T3	28-09-2001
			HU 76785 A2	28-11-1997
			JP 10506624 T	30-06-1998
			NO 971342 A	21-03-1997
			NZ 293249 A	29-04-1999
			PT 783505 T	30-08-2001
			US 5869485 A	09-02-1999
<hr/>				
WO 9716452	A	09-05-1997	AU 7296896 A	22-05-1997
			BR 9611157 A	30-03-1999
			CA 2234609 A1	09-05-1997
			CN 1202896 A ,B	23-12-1998
			WO 9716452 A1	09-05-1997
			EP 0874846 A1	04-11-1998
			JP 11514336 T	07-12-1999
			ZA 9609168 A	01-05-1997
<hr/>				